

## *Nigrospora Sphaerica* Products from the Flowering Dogwood Exhibit Antitumorigenic Effects Via the Translational Regulator, Ps6 Ribosomal Protein

Asha Maheshwari<sup>1,2</sup>, Margaret Mmbaga<sup>1</sup> & Quincy Quick, A.<sup>2\*</sup>

<sup>1</sup>Department of Agricultural and Environmental Sciences, Tennessee State University, Nashville, TN, USA

<sup>2</sup>Department of Biological Sciences, Tennessee State University, Nashville, TN, USA

\***Correspondence to:** Dr. Quincy Quick, A., Department of Biological Sciences, Tennessee State University, 3500 John A. Merritt Blvd, Nashville, TN, 37209 USA.

### Copyright

© 2018 Dr. Quincy Quick, A., *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 01 June 2018

Published: 13 June 2018

**Keywords:** *Natural Product; Fungi; Secondary Metabolite; Intracellular Signaling; Cancer*

### Abstract

The utility of diverse species of endophytic fungi as a viable source for drug agents with clinical applicability for the treatment of human diseases continues to expand. In this study we examined secondary metabolites of *Nigrospora sphaerica* isolated from the flowering dogwood, *Cornus florida* L., for their anticancer properties on lung cancer and glioblastoma. Molecular identification of *N. sphaerica* was determined using ITS-rDNA sequence. The expression of translational pathway proteins was examined after exposure to various crude extract concentrations (2µg/ml, 4µg/ml, 8µg/ml) using immunoblotting procedures, while tumor cell migration analysis was performed using boyden chamber assays. Crude *N. sphaerica* extracts exhibited antiproliferative and antimigratory effects on solid tumors as determined by cell proliferation and cell migration assays, respectively. The antitumorigenic effects of *N. sphaerica* were as a consequence of negatively regulating the PI3K/Akt/mTOR translational control signaling pathway, a canonical mechanistic axis that contributes to the maintenance and progression of several human cancers. To our knowledge this is the first

evidence that demonstrates *N. sphaerica* from *C. florida* inhibits tumor cell migration, and thus disease recurrence a major factor in the therapeutic resistance of cancers to chemotherapeutic agents.

## Background

Endophytic fungi are organisms that colonize the internal tissues of plants without damaging the host [1], and have proven to be a robust source for the discovery of novel secondary metabolites that exhibit a wide range of biological and biomedical applications [2-4]. A candidate source and potential host of endophytic fungi with biomedical applications is the flowering dogwood, *Cornus florida* L., which has been shown to produce antiparasitic compounds for the treatment of malaria [5]. Germane to the present study *C. florida* has also been shown by Vareed *et al.*, [6] to produce the phenolic pigment anthocyanin, which displayed antitumor activity in colon, breast, lung, stomach, and central nervous system cancers. To this end the current study examined the antitumor activity of crude extracts prepared from the endophytic fungus *Nigrospora sphaerica* isolated from *C. florida*, which has previously been described as an endophyte resource for several biomedical applicable compounds such as antibiotics [7-8], antiviral anthraquinones and azaphilones [9], and the antifungal griseofulvin [10]. Of specific significance the alkaloid chemotherapeutic drug, vinblastine, is also a product of the endophytic fungus *N. sphaerica* albeit isolated from *Catharanthus roseus* (Madagascar periwinkle) [11]. However, the clinical caveat with the utility of the microtubule poison vinblastine for the treatment of human cancers has been its associated toxicities, making it imperative to identify sources for chemotherapeutic agents with improved toxicity profiles. Here we examined the antiproliferative effect of crude extracts from *N. sphaerica* on lung and glioblastoma cancer cell lines, as well as the ability of *N. sphaerica* extracts to inhibit cancer cell migration. The results of this study provide experimental evidence that the endophytic fungus *N. sphaerica* in flowering dogwoods is a promising source of bioactive metabolites with potential antitumor and antimetastasis activities.

## Methods

### Sample collection and Fungal isolation

Healthy stem samples were collected from mature *C. florida* plants at Otis L. Floyd Nursery Research Centre (Tennessee State University, McMinnville, TN). The samples were placed in sterile bags, stored at 4°C and processed within 24 hours of sampling. Endophytic fungi were isolated using methods as previously described by Schulz *et al.*, [12] with some modification. Stem samples were cleaned thoroughly under running tap water and cut into 1.0-1.5 cm pieces. Samples were next surface sterilized with 70% ethanol for 1 minute and 10% sodium hypochlorite for 3-5 minutes. Samples were subsequently rinsed three times with sterile distilled water, blotted on paper towel and allow to air dry. Next five segments of vascular tissue (3-4mm) were transferred onto petri dishes containing acidified potato dextrose agar, incubated at room temperature and observed for appearance of mycelial growth. Pure fungus was obtained by at least two or three successive sub-culturing on potato dextrose agar (PDA; Sigma-Aldrich, St. Louis, MO).

## Molecular identification and phylogenetic analysis

Fungal genomic DNA was extracted using the FastDNA kit (MP Biomedicals, Santa Ana, CA) per manufacturers' instructions. Next PCR of the Internal Transcribed Spacer (ITS) region was performed using the universal primers ITS1 and ITS4 [13]. Subsequently, the PCR product was purified using Exosap (USB-Affimetrix, Santa Clara, CA), and sequenced by Eurofin Genomics (Louisville, KY). Next the ITS sequence was analyzed using BLASTN (GenBank National Centre of Biotechnology Information-NCBI; <https://www.ncbi.nlm.nih.gov>) and the identity of the organism was determined based on the closest match in the GenBank database. DNA sequence similarities of  $\geq 99$ -100% homology were used to identify the fungus *N. sphaerica* studied here. Subsequently, the *N. sphaerica* ITS sequence and reference sequences retrieved from GenBank were aligned using ClustalW prior to phylogenetic tree construction in MEGA7 software [14]. The neighbor-joining (NJ) method was used to infer evolutionary history [15]; the p-distance method was used to compute evolutionary distances [16]; and bootstrap analysis with 1000 replication was used to assess the robustness of the phylogenetic tree [17].

## Crude Extract preparation

For crude extract preparation 5-7, 6mm<sup>2</sup> agar plugs of fungal culture were inoculated in 1000 ml erlenmeyer flasks containing 300 ml of potato dextrose broth (Sigma Aldrich, St Louis, MO; USA). Flasks were then incubated at 28°C on a rotary shaker at 160 rpm for 10 days. After incubation, the fermented medium was centrifuged at 4000 rpm to separate mycelia from the supernatant. Subsequently, supernatant was filter sterilized (0.22 µm) and extracted three times with equal volumes of ethyl acetate (Fisher Scientific, Fair Lawn, NJ; USA) using separatory funnel. Next the organic phase was removed using rotary evaporator under vacuum and resultant crude extract was lyophilized. The stock solution of 2.0 mg/ml crude extract was prepared in dimethyl sulfoxide (DMSO; Amresco Solon, OH; USA) and stored at -20°C for further analysis.

## Cells culture conditions and reagents

U251 human glioblastoma cells were purchased from Sigma-Aldrich (St. Louis, MO; USA) and A549 human lung carcinoma cells were obtained from the American Type Culture Collection (Manassas, VA; USA). Both cell lines were maintained in Dulbecco's Modified Eagles Medium-DMEM (Invitrogen, Carlsbad, CA; USA) supplemented with 10% Fetal Bovine Serum (Invitrogen, Carlsbad, CA; USA), 2mM L-glutamine (Invitrogen, Carlsbad, CA; USA), 100nM MEM non-essential amino acids (Invitrogen, Carlsbad, CA; USA), and penicillin-streptomycin (Invitrogen, Carlsbad, CA; USA) at 37°C and 5% CO<sub>2</sub>.

## Crystal violet cell proliferation assay

U251 and A549 cells were plated in 24 well plates and treated with 2 µg/ml, 4 µg/ml, 8 µg/ml, and 10 µg/ml of fungal crude extract, while vehicle treated control cells were treated with DMSO. For dose response experiments cells were treated with *N. sphaerica* (2 µg/ml, 4 µg/ml, 8 µg/ml, and 10 µg/ml) fungal extract or vehicle and allowed to incubate for 48h, while for time course experiment cells were plated, treated as explained above and incubated for 2, 4 and 6 days. At the end of each time point the tissue culture medium

was removed, the cell monolayer was fixed with 100% methanol for 5 minutes and stained with 0.5% crystal violet in 25% methanol for 10 minutes. Cells were then washed three times five minutes each with distilled water to remove excess dye and allowed to dry overnight at room temperature. The incorporated dye was then solubilized in 0.1M sodium citrate (Sigma-Aldrich St. Louis, MO; USA) in 50% ethanol. Next, 100µl of experimental samples were transferred to 96 well plates and optical densities were recorded at 595 nm using an X-mark microplate absorbance spectrophotometer (BioRad, Hercules, CA; USA).

### **Cell migration**

Motility assays were conducted according to manufacturer's instructions (CellBiolabs Inc, San Diego, CA; USA). Cell suspensions containing  $0.5-1.0 \times 10^6$  cells/ml were treated with DMSO (vehicle control) or 2 µg/ml of crude extract in serum free media, while 500 µl of media containing 10% fetal bovine serum was added to the lower chamber of the migration plate. Next 300 µl of cell suspension containing vehicle control or 2 µg/ml of crude extract were then added to the inside of each insert and allowed to incubate for 24 hours at 37°C and 5% CO<sub>2</sub>. Subsequently non-migratory cells were removed from plate inserts (per manufacturer's instructions), while migratory cells were stained with crystal violet and the dye solubilized (per manufacturer's instructions). Optical densities were read at 595nm using an X-mark microplate absorbance spectrophotometer (BioRad, Hercules, CA-USA).

### **Caspase 3 activity**

To assess caspase 3 activity cells were plated in serum-free DMEM for 24 h, treated with 2 µg/ml, 4 µg/ml, 8 µg/ml crude extract, or vehicle (DMSO) and allowed to incubate for 24 h. After incubation, cells were rinsed with cold phosphate buffer saline (PBS; Bio-Rad Laboratories, Hercules, CA, USA) and lysed in CelLytic M cell lysis reagent (Sigma-Aldrich, St Louis, MO, USA). Protein concentrations were then determined using the Bradford method. Next, caspase 3 activity was determined using the CaspACE Assay System (Promega, Madison, WI, USA) according to the manufacturer's instructions.

### **Western Blotting**


For western blot experiments cells wells were treated with 2 µg/ml, 4 µg/ml, 8 µg/ml crude extract or vehicle (DMSO) for three hours, rinsed with PBS, and lysed with CelLytic M cell lysis reagent (Sigma-Aldrich, St Louis, MO; USA). Protein concentrations were subsequently determined using the Bradford method. Proteins were next run in 4-15% SDS- polyacrylamide gels (BioRad, Hercules, CA; USA) and transferred to nitrocellulose membranes. Nitrocellulose membranes were then incubated overnight at 4°C with primary antibodies to detect the protein expression of pAkt, pS6 ribosomal protein, p90 RSK, 4EBP1, GAPDH, and β actin (CellSignaling Technology, Danvers, MA; USA). Nitrocellulose membranes were next washed three times 5 minutes each in PBS containing 0.05% Tween 20, incubated with an HRP-conjugated secondary antibody for 3h at room temperature and washed. Proteins were visualized and analyzed by enhanced chemiluminescence (Thermo Scientific, Nashville, TN; USA) and a UVP BioSpectrum imaging system (UVP, Upland, CA; USA).

## Statistical analysis

Cell proliferation, migration, and time-course experiments were each performed at least three times in duplicate. Means were determined by averaging duplicate samples within each independent experiment. Students t-tests and ANOVA with Sidak's posthoc statistical analysis were used to evaluate significance.

## Result

The endophyte *N. sphaerica* has been previously reported in a number of plant species [11, 18 - 20]. To our knowledge this study is the first report to show the association of *N. sphaerica* with *C. florida* as confirmed using ITS-rDNA sequence and phylogenetic analysis (Figure 1).




**Figure 1.** Neighbor-joining phylogenetic tree generated by *p*-distance method based on ITS sequence of endophytic *N. sphaerica* (▲) isolated from *C. florida*. The optimal tree with the sum of branch length = 0.02020992 is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches.


Additionally, few studies have evaluated the anticancer properties of secondary metabolites from *N. sphaerica*. Initial experimental studies on the effect of the endophytic fungus *N. sphaerica* on human cancer cells was determined via dose response analysis on lung cancer and glioblastoma cell lines, which displayed a concentration dependent effect (Figures 2 and 3). Glioblastoma cells treated with 2 µg/ml of *N. sphaerica* crude extract showed a 25% decrease in cell viability and an 84% decrease in cell viability when treated with 10 µg/ml, while lung cancer cells treated with these concentrations displayed a 17% and 62% decrease, respectively, as compared to vehicle treated control cells (Figure 2). The quantitative effect of this *Nigrospora* fungus on lung and glioblastoma cancer cell viability was further supported by ANOVA analysis which



showed a statistically significant difference ( $p < 0.05$ ) across all concentrations examined. Subsequently, time course analysis were performed on lung and glioblastoma cancer cells exposed to crude extracts of *N. sphaerica*.




**Figure 2.** Dose response effects of *N. sphaerica* on lung cancer (A549) and glioblastoma cells (U251). Cells treated with 2 µg/ml – 10 µg/ml of *N. sphaerica* crude extract caused a reduction in cell viability. Data shown is representative of at least four experiments performed in duplicate (means ± SE) that showed similar results (\*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$  compared to vehicle treated control cells).



**Figure 3.** Qualitative dose response assessment of *N. sphaerica* crude extract on lung cancer (A-E) and glioblastoma cells (F-J). Concentrations of *N. sphaerica* crude extract: DMSO vehicle control (A, F); 2 µg/ml (B, G); 4 µg/ml (C, H); 8 µg/ml (D, I); 10 µg/ml (E, J). Shown is a single experiment representative of four independent experiments that displayed similar results.


Consistent with dose response experiments cells treated with *N. sphaerica* crude extracts ranging in concentration between 2  $\mu\text{g/ml}$  and 10  $\mu\text{g/ml}$  also displayed concentration dependent effects of crude extracts on glioblastoma and lung cancer cells, with the highest doses having the most pronounced effect on decreasing cell viability (Figure 4). This is supported by a 93% and 96% decrease in viable lung cancer and glioblastoma cells, respectively, observed 6 days' post-exposure with 10  $\mu\text{g/ml}$  of *N. sphaerica* crude extract, as compared to vehicle treated control cells examined at this same time point (Figure 4). Additionally, treatment with 10  $\mu\text{g/ml}$  of *N. sphaerica* induced a cytostatic cellular response in lung cancer cells and a cytotoxic cellular response in glioblastoma cells over the six-day time period (Figure 4). These data are consistent with the antiproliferative effects of *N. sphaerica* products observed in leukemia and breast cancer cells [11, 21].




**Figure 4.** Time course analysis of *N. sphaerica* crude extract on lung cancer (A549) and glioblastoma cells (U251). *N. sphaerica* crude extract (10  $\mu\text{g/ml}$ ) induced a cytotoxic and cytostatic cellular response as compared to vehicle treated control cells. Shown is an experiment representative of three independent experiments performed in duplicate (means  $\pm$  SE) that displayed similar results.

To gain insight into the role that apoptotic cell death played in the reductions in cancer cell viability as a consequence of exposure to *N. sphaerica* crude extracts, caspase 3, a known executioner of apoptosis, was evaluated in lung cancer and glioblastoma cells (Figure 5) post-exposure to crude extracts. Our data showed a diminutive increase in caspase 3 activity in both cell lines treated with *N. sphaerica* crude extracts (Figure 5). Additionally, we assessed the alternative cell death mechanism autophagy, as a means that underlie reduced cell viability of lung and glioblastoma cells treated with *N. sphaerica* crude extract. Immunoblotting data showed no change in the autophagy marker LC3 in response to *N. sphaerica* crude extract exposure (data not shown). We further analyzed the expression of proteins that regulate translational control (pAkt, pS6 ribosomal protein, 4EBP1) and ERK (p90RSK) signaling pathways as mechanistic contributors of the anti-proliferative effect of *N. sphaerica* crude extract on solid tumor cell lines studied here. *N. sphaerica* crude extract exposure caused a demonstrative decrease in the expression of the translational regulator pS6

ribosomal protein in lung cancer cells (Figure 6) but no measurable change in the ERK signaling protein p90RSK. In contrast *N. sphaerica* crude extract had no effect on translational control and ERK signaling proteins evaluated here in glioblastoma cells (data not shown).




**Figure 5.** Evaluation of apoptosis in response to *N. sphaerica* crude extract. Treatment of lung cancer (A549) and glioblastoma (U251) cells with *N. sphaerica* crude extract induced a slight increase in caspase 3 activity. Caspase 3 activity data shown is representative of at least three experiments performed in duplicate (means ± SE) that showed equivalent results, while immunoblots presented are representative of at least three experiments that showed similar results.





**Figure 6.** Western blot analysis of translational control and ERK1/2 signaling mediators in lung cancer cells. *N. sphaerica* crude extract promoted a down-regulation of pS6 ribosomal protein in A549 lung cancer cells. Immunoblots presented are representative of at least four experiments that showed equivalent results. GAPDH and actin were used as loading controls to assess that lanes were loaded with the same amount of total proteins.

Because tumor recurrence is a manifestation of the resistance of many cancers to clinically used chemotherapeutic agents, we next examined the efficacy of *N. sphaerica* crude extracts to inhibit lung cancer and glioblastoma cell migration. Boyden chamber assays revealed that cells treated with 2  $\mu\text{g}/\text{ml}$  of *N. sphaerica* crude extracts decreased ( $p < 0.05$ ) tumor cell migration as compared to vehicle treated control cells (Figure 7), paralleling recent findings by Mady *et al.*, [22] that showed the endophytic fungus *Penicillium chrysogenum* impaired breast cancer cell migration.



**Figure 7.** *N. sphaerica* crude extract impairs solid tumor cell migration. Lung cancer (A549; left panel) and glioblastoma cell (U251; right panel) migration were significantly diminished after treatment with 2  $\mu\text{g}/\text{ml}$  of *N. sphaerica* crude extract. Data shown are representative of three independent experiments performed in duplicate (means  $\pm$  SE) showing similar results (\*  $p < 0.05$  compared to vehicle treated control cells). Vehicle treated control cells (blue bars); cells treated with 2  $\mu\text{g}/\text{ml}$  of *N. sphaerica* crude extract for 24 hours (red bars).

## Discussion

We have shown here that metabolic products of the endophytic fungus *N. sphaerica* has antitumorigenic properties as exhibited by the antiproliferative effects of *N. sphaerica* crude extracts from *C. florida* on lung cancer and glioblastoma cells. Our findings are consistent with previous experimental studies that have also shown secondary metabolites produced by *N. sphaerica* isolated from *C. roseus* and *Asteraceae* inhibited the proliferation of breast cancer and leukemia cells, respectively. More specifically Ayob *et al.*, [11] demonstrated that the vinca alkaloid, vinblastine produced by *N. sphaerica* from *C. roseus* inhibited breast cancer cell growth. It should be mentioned that the concentration however of vinblastine used to inhibit breast cancer cells

by Ayob *et al.*, [11] was three times that of the highest concentration of *N. sphaerica* crude extract used in this study to inhibit the growth of solid cancer cell lines. This suggests that cancer cells may be more sensitive to *N. sphaerica* products extracted from *C. florida* and that endophytic fungi products from this plant are more effective at inhibiting cancer cell proliferation. Additionally, Gallo *et al.*, [21] showed that diterpene aphidicolin, a *N. sphaerica* product of *Asteraceae* had an antiproliferative effect on leukemia cells that was attributed in part to G2/M cell cycle arrest, apoptosis, and autophagy. In contrast *N. sphaerica* crude extracts in our study did not induce apoptotic cell death or an autophagic response, but instead promoted a decrease in the translational regulator, pS6 ribosomal protein an effector molecule in the PI3K/Akt/mTOR signaling pathway regulated by mTORC1 (Raptor-mTOR) a known regulator of cell growth, glucose and lipid metabolism, autophagy, and protein synthesis [23]. Although the change in pS6 ribosomal protein expression provides a mechanistic basis for the antiproliferative effects of lung cancer and glioblastoma cells seen in response to *N. sphaerica* crude extracts, the downregulation of this translational regulatory protein has also been associated with impeding cancer cell migration. In a study by Kim *et al.*, [24] it was demonstrated that genetic downregulation of pS6 ribosomal protein inhibited esophageal cancer cell migration and invasion. This was further supported in studies by Chen *et al.*, [25] who observed this same phenomenon in non-small cell lung cancer. Taken together these investigations support the notion that impairment of lung cancer and glioblastoma cell migration to *N. sphaerica* crude extracts isolated from *C. florida* is also a consequence of pS6 ribosomal protein reduction.

## Conclusion

The anti-migratory effects in conjunction with the antiproliferative properties of *N. sphaerica* crude extracts from *C. florida* on solid tumor cell lines observed in our study provide evidence that endophytic fungi isolated from this plant is able to impede the continued development and recurrence of human cancers. Furthermore, findings here expand the pool of botanical resources that can be used to evaluate secondary metabolites from endophytic fungi for their anticancer properties to include the flowering dogwood, *C. florida*.

## Acknowledgements

This work was partly supported by United States Department of Agriculture, National Institute of Food and Agriculture: Award Number 2010-38821-21477. The authors declare that they have no competing interests.

## Bibliography

1. Petrini, O. (1991). Fungal Endophytes of Tree Leaves. In: Andrews, J. H., Hirano, S. S. *Microbial Ecology of Leaves*. New York (NY): Springer Verlag.
2. Strobel, G. A. (2002). Rainforest endophytes and bioactive products. *Crit Rev Biotechnol.*, 22(4), 315-33.
3. Nair, D. N. & Padmavathy, S. (2014). Impact of Endophytic Microorganisms on Plants, Environment and Humans. *The Scientific World Journal*.
4. Venugopalan, A. & Srivastava, S. (2015). Endophytes as in vitro production platforms of high value plant secondary metabolites. *Biotechnology Advances*, 33(6), 873-887.

---

Quincy Quick, A. (2018). *Nigrospora Sphaerica* Products from the Flowering Dogwood Exhibit Antitumorigenic Effects Via the Translational Regulator, Ps6 Ribosomal Protein. *CPQ Cancer*, 1(1), 01-12.

5. Graziose, R., Rojas-Silva, P., Rathinasabapathy, T., *et al.* (2012). Antiparasitic compounds from *Cornus florida* L. with activities against *Plasmodium falciparum* and *Leishmania tarentolae*. *J Ethnopharmacol.*, 142(2), 456-61.
6. Vareed, S. K., Reddy, M. K., Schutzki, R. E., *et al.* (2006). Anthocyanins in *Cornus alternifolia*, *Cornus controversa*, *Cornus kousa* and *Cornus florida* fruits with health benefits. *Life Sci.*, 78(7), 777-84.
7. Yang, K. L., Wei, M. Y., Shao, C. H., *et al.* (2012). Antibacterial anthraquinone derivatives from a Sea Anemone derived fungus *Nigrospora sp.* *J. Nat. Prod.*, 75(5), 935-941.
8. Ibrahim, D., Lee, C. C., Yenn, T. W., *et al.* (2015). Effect of the extract of endophytic fungus, *Nigrospora sphaerica* CL-OP 30, against the growth of methicillin- resistant *Staphylococcus aureus* (MRSA) and *Klebsiella pneumonia cells*. *Tropical Journal of Pharmaceutical Research*, 14(11), 2091-2097.
9. Zhang, S. P., Huang, R., Li, F. F., *et al.* (2016). Antiviral anthraquinones and azaphilones produced by an endophytic fungus *Nigrospora sp.* from *Aconitum carmichaeli*. *Fitoterapia*, 112, 85-9.
10. Zhao, J. H., Zhang, Y. L., Wang, L. W., *et al.* (2012). Bioactive secondary metabolites from *Nigrospora sp.* LLGLM003, an endophytic fungus of the medicinal plant *Moringa oleifera* Lam. *World J Microbiol Biotechnol*, 28(5), 2107-12.
11. Ayob, F. W., Simarani, K., Zainal Abidin, N., *et al.* (2017). First report on a novel *Nigrospora sphaerica* isolated from *Catharanthus roseus* plant with anticarcinogenic properties. *Microb Biotechnol.*, 10(4), 926-32.
12. Schulz, B., Wanke, U., Draeger, S., *et al.* (1993). Endophytes from herbaceous plants and shrubs: effectiveness of surface sterilization methods. *Mycol Res.*, 97(12), 1447-1450.
13. White, T. J., Bruns, T., Lee, S., *et al.* (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*, New York (NY): Academic Press, Inc.
14. Kumar, S., Stecher, G., Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33(7), 1870-1874.
15. Saitou, N. & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular Biology and Evolution*, 4(4), 406-425.
16. Nei, M. & Kumar, S. (2000). *Molecular Evolution and Phylogenetics*. New York (NY): Oxford University Press.
17. Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39(4), 783-791.

18. Gallo, M. B., Chagas, F. O., Almeida, M. O., *et al.* (2009). Endophytic fungi found in association with *Smallanthus sonchifolius* (Asteraceae) as resourceful producers of cytotoxic bioactive natural products. *J Basic Microbiol.*, 49(2), 142-51.
19. Metwaly, A. M., Kadryb, H. A., El-Helab, A. A., *et al.* (2014). Nigrosphaerin A a new isochromene derivative from the endophytic fungus *Nigrospora sphaerica*. *Phytochem Lett.*, 7, 1-5.
20. Santos, I. P., Silva, L. C. N., Silva, M. V., *et al.* (2015). Antibacterial activity of endophytic fungi from leaves of *Indigofera suffruticosa* Miller (Fabaceae). *Front. Microbiol.*, 6, 350.
21. Gallo, M. B. C., Falso, M. J. S., Balem, F., *et al.* (2014). The Anti-Promyelocytic leukemia mode of action of two endophytic secondary metabolites unveiled by a proteomic approach. *Planta Med.*, 80(6), 473-481.
22. Mady, M. S., Mohyeldin, M. M., Ebrahim, H. Y., *et al.* (2016). The indole alkaloid Meleagrins, from the olive tree endophytic fungus *Penicillium chrysogenum*, as a novel lead for the control of c-Met-dependent breast cancer proliferation, migration and invasion. *Bioorg Med Chem.*, 24(2), 113-22.
23. Liu, Y., Wan, W. Z., Li, Y., *et al.* (2017). Recent development of ATP-competitive small molecule phosphatidylinositol-3-kinase inhibitors as anticancer agents. *Oncotarget.*, 8(4), 7181-7200.
24. Kim, S. H., Jang, Y. H., Chau, G. C., Pyo, S. & Um, S. H. (2013). Prognostic significance and function of phosphorylated ribosomal protein S6 in esophageal squamous cell carcinoma. *Mod Pathol.*, 26(3), 327-35.
25. Chen, B., Tan, Z., Gao, J., Wu, W., Liu, L., Jin, W., Cao, Y., Zhao, S., Zhang, W., Qiu, Z., *et al.* (2015). Hyperphosphorylation of ribosomal protein S6 predicts unfavorable clinical survival in non-small cell lung cancer. *J Exp Clin Cancer Res.*, 34, 126.