

Proximate, Micronutrient, Functional and Microbial Properties of Complementary Foods Sold in Three Localities in South West Region of Cameroon

Arrey Oben Ebob Ashu¹, Tiepma Ngongang Eurydice Flore¹, Bernard Tiencheu^{*}, Aduni Ufuan Achidi¹, Mbame Efeti Claudine¹, Noel Tenyang² & Feumba Dibamda Romelle¹

¹*Department of Biochemistry and Molecular Biology, Faculty of Science, University of Buea, Buea, Cameroon*

²*Department of Biological Science, Faculty of Science, University of Maroua, Maroua, Cameroon*

***Correspondence to:** Dr. Bernard Tiencheu, Department of Biochemistry and Molecular Biology, Faculty of Science, University of Buea, Buea, Cameroon.

Copyright

© 2021 Dr. Bernard Tiencheu, *et al.* This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: 08 December 2021

Published: 27 December 2021

Keywords: *Complementary Food; Nutrient Composition; Adequacy; Children; Reference Standard*

Abstract

Mothers and caregivers typically feed infants according to their culture, purchase power and level of awareness with no due diligence to nutritional quality of the diet. Scientific evidence on nutritional adequacy of predominant complementary foods is critical for planning and prioritizing interventions. The purpose of the current study was to evaluate the quality of complementary foods. Some complementary food types Vita Force (VT), Soytime (ST), Soyaconia (SC), Soya Pap (SP), Tanty Reine (TR), Phospatine, Dina Baby (DB), were assayed for proximate composition, energy and micronutrients using standard method of AOAC, (2005) [1]. Adequacy of the complementary foods in nutrients for complementary feeding purposes was assessed as a ratio between actual composition and recommended composition of complementary foods. Microbiological analysis was done using standard methods of Olurunjuwon *et al.*, (2014) [2]. Data obtained was statistically analyzed using GraphPad Instat 3. The results for the proximate analysis (carbohydrate, protein, fat,

ash, fibre) of the complementary food samples were comparatively lower than the recommendation by FAO (2006) [3]. Micronutrient content of the weaning foods as well as the functional properties was significantly different from that of the standard. Mineral composition showed that none of the different weaning foods had the recommended value for the minerals Ca, Na, K, Fe, Zn, Mg except for one of the semi-scaled CF Tanty which had the right amount (48.6%) of Mg mineral. Microbiological load for weaning foods determines the wholesomeness of weaning foods for consumption as the Coliform counts were in acceptable range (less than 3cfu/g) but homemade food samples like kwacoco and pap had higher value for total bacterial count of 200cfu/g. Hence, most of the weaning foods sold in Fako, South West region of Cameroon are not adequate for children's normal nutrition and growth.

Introduction

Exclusively, breast feeding for the first 6 months is the world health organization (WHO)'s recommended method of feeding full-term infants by healthy, well-nourished mothers. However, after 6 months' breast milk alone is not sufficient both in quantity and in quality to meet the nutritional requirements of the child (Kalanda *et al.*, 2006). Adequate foods known as complementary foods need to be introduced, which begins with liquid foods and gently move up to solid foods; this is the weaning process. Weaning is when foods other than breast milk are given to infant's diet while gradually reducing breastfeeding (WHO, 2014). The introduction of weaning foods is often accompanied by stress and ill health for infants in developing countries mostly because the foods are not properly tailored to the infant's needs [4]. Many traditional weaning foods in Africa are only a slight modification of adult food involving only mashing and dilution without taking into consideration the special nutritional requirements of young children. Adults diet especially in developing countries, consist of highly starchy staples and unless properly modified, unsuitable for infants and young children with their small gastric capacities [5]. However, during the process of complementary feeding, children need foods that are not strong, hygienic, energy and nutrient-dense to reach their high nutritional requirement (WHO, 2013). Complementary foods in third world countries are mostly prepared under non sterile environments using water from un-hygienic sources thus making the child vulnerable to weaning diarrhea which causes a dilemma to both mother and infant; to wean or not to wean, which is termed the weaning's dilemma [6]. Therefore, the aim of this work which is to evaluate the adequacy of complementary foods sold in the South West region of Cameroon.

Material and Methods

Some locally produced baby foods like Sweet potato custard, Banana custard, Soyaconia (SC), Bionutri, Tanty Reine (TR) and Dina Baby (DB) were purchased from local stores in Buea, Mutengene and Limbe in the South West Region of Cameroon. A standard (Phosphatine) a commercially sold baby food was obtained from a super market and used as a control. The samples were packed and sealed in polyethylene bags at ambient temperature ($26 \pm 2^\circ\text{C}$), transported directly to the laboratory stored in the refrigerator until further analysis.

Proximate Composition, Vitamins and Minerals were carried out following the method of [1].

Microbial Analysis

The yeast, coliform and bacterial loads were analyzed using the Sabouraud dextrose agar, violet red bile lactose agar and plate count agar media respectively [2].

Total Yeast Count

Sabouraud Dextrose agar (SDA) (4.2g) was dissolved in 100ml of distilled water. It was autoclaved at 121°C for 1 hour in an electric pressure steam sterilizer (model No.25X) and allowed to cool to about 45°C. The work environment was sterilized with 70% alcohol. 1ml of each sample was diluted in 9ml of water and 1ml of the diluted samples poured into labeled petri-dishes with the use of a micropipette (Gilson Pipetman, 060087N). Then 10ml of the medium (SDA) was put into a petri-dish and shaken gently to mix with the sample. This solidified and formed a gel in the petri dishes. They were then incubated (DHP-9050) at 37°C for 24 hours. After 24 hours, the total yeasts were counted as colony forming units (CFU) [2].

Total Bacterial Count

Plate count agar (PCA) (2.4g) was diluted in 100ml of distilled water. It is autoclaved at 121°C for 1 hour in a sterilizer (model No.25X) and left to cool to about 45°C. The working space was sterilized with 70% alcohol. For every sample, 1ml was diluted in 9ml of water and 1ml of the diluted samples poured into labeled petri-dishes using a micropipette (Gilson Pipetman, 060087N). Then 10ml of the medium (PCA) was put into the petri dish and shaken gently to mix with the sample. This molded and produced a gel in the petri dish. It was then incubated (DHP-9050) at 37°C for 24 hours. After 24 hours, the total bacteria load was expressed as colony forming units (CFU) [2].

Total Coliform Count

Violet red agar (6.2g) was dissolved in 150ml of distilled water. This solution was allowed to boil while shaking over a bunsen burner flame until it was completely dissolved. It was then allowed to cool to about 45°C. The work environment is sterilized with 70% alcohol. For each sample, 1ml was diluted in 9ml of water and 1ml of the diluted samples poured into labeled petri-dishes with the use of a micropipette (Gilson Pipetman, 060087N). Then 10ml of the medium (SDA) was poured into the petri dish and agitated gently to homogenize with the sample. This solidified and formed a gel in the petri dish. It is then incubated (DHP-9050) at 42°C for 24 hours. After 24 hours, the total coliforms were counted as colony forming units (CFU) [2].

Functional Properties

Oil Absorption Capacity (OAC) and Water Absorption Capacity (WAC):

AOAC and WAC were determined using the method described by Lin *et al.* (1974). One gram of protein sample was vortexed with 10ml of sunflower oil or distilled water respectively for 30s. The emulsions were

incubated at room temperature (20°C) for 30 min and then centrifuged at 13,600g for 10min at 25°C. The supernatant was decanted and drained at 45°C for 20 min.

$$\text{Oil absorption capacity} = \frac{\text{vol of oil or water absorbed}}{\text{weight of protein sample}}$$

Determination of the Density

Bulk density was calculated using the method of Murphy *et al.*, (2002) for 5mins to the sample was put into a 10ml graduated cylinder up to the 10ml mark. The cylinder was rapped to eliminate air space between the particles of the blends in the cylinder. The density in g/cm³ was calculated as the weight (g) of flour divided by the volume (cm³) of flour [7].

Dispersibility

It was estimated following the method described by Kulkarni *et al.*, (1991) [8]. 1g of sample was weighed and mixed with 10ml of water and stirred. The mixture was then shaken vigorously and allowed to settle for 3hours. Then to obtain the dispersibility, the volume of particles which had settled was subtracted from 10ml and reported as a percentage. %Dispersibility = 10ml -Volume of particles which settled.

Foaming Capacity (FC):

The foaming capacity and stability were determined according to the method described by Narayana and Narsinga (1982) [9] with some changes. The 1.0g flour sample was added to 50ml of distilled water at 30°C using a thermometer in a graduated cylinder. The suspension was mixed and agitated for 5min to foam. After agitation, the foam volume was calculated as foam capacity.

$$\text{Foam capacity (\%)} = \frac{(\text{volume of foam AA} - \text{volume of foam BA}) * 100}{\text{volume of foam BA}}$$

where, AA= after agitation, BA= before agitation. The volume of foam was recorded 1 hour after whipping to determine foam stability as per percent of initial foam volume

Swelling Index

Abbey and Ibeh (1987) methods were followed. 1g of the flour sample was introduced into a 10ml graduated cylinder. 5ml of water which has been distilled was gently added and the volume occupied by the sample noted. The sample was kept standing for some time and was soaked in water for 1 hour and the volume occupied after swelling was measured.

$$\text{Swelling Index} = \frac{\text{volume occupied by sample after swelling}}{\text{volume occupied by sample before swelling}}$$

Protein Solubility

Protein solubility was tested in the pH ranges 1 to 10. One gram of each sample was suspended in 5ml distilled water and the pH of the suspension was adjusted to a specific value (1-10) using 0.1N HCL or NaOH solutions. These suspensions were agitated over a shaker for one hour at room temperature, the pH was checked and readjusted in the range of 1 to 10, then centrifuged at 1500rpm for 15 minutes. The soluble protein was determined by the method of biuret at 540nm (Rosebrough *et al.*, 1951) using BSA as standard. Protein solubility was expressed as the percentage of the total protein of the original sample that was present in the soluble fraction.

Statistical Analysis

Data collected was analyzed using the IBM-Statistical Package for Social Sciences (IBM-SPSS) version 20. Continuous variables were summarized into means and standard deviations (SD) and categorical variables reported as frequencies and percentages were used to evaluate the descriptive statistics. The differences in proportions were evaluated using Pearson's Chi-Square (χ^2). Significant levels were measured at 95% confidence interval (CI) with significant differences set at <0.05. Statistical analysis of nutritional composition was done using GraphPad InStat 3 (trial) and the differences between the mean values were evaluated at $P < 0.05$ using Duncan's multiple range test.

Results

Nutritional Evaluation

The result of proximate composition, β carotene and vitamin C are presented (Table 1 and Table 2 respectively).

Table 1: Proximate analysis of complementary foods

| Samples | Moisture (%) | Fat (%) | Fibre (%) | Protein (%) | Carbohydrate (%) | Ash (%) | Energy Kcal/100g |
|--------------------------------|--------------|------------|------------|-------------|------------------|-----------|------------------|
| WHO/FAO Reference pattern 2006 | 5 | 8 | 3.8 | 15 | 64-68 | 2.9 | 400 |
| Semiscaled CF | | | | | | | |
| Soyaconia | 4.6±0.84* | 12.95±0.4* | 10.2±0.84* | 17.30±1.82 | 58.70±0.71* | 4.25±0.35 | 420.55 |
| Tantyreine | 8.62±0.10* | 21.56±0.2* | 13.7±0.98* | 12.08±0.62* | 54.85±0.49* | 4.15±2.19 | 461.22 |

| | | | | | | | |
|----------------------|-------------|-------------|------------|-------------|-------------|-----------|--------|
| Banana custard | 11.15±1.40* | 10.73±1.4* | 12.3±0.14* | 12.95±2.47 | 59.65±1.63 | 4.5±0.14 | 386.97 |
| Sweet potato custard | 8.5±0.47* | 10.51±0.8* | 13.1±0.98* | 11.42±2.16* | 64.45±1.34 | 4.4±0.28 | 398.07 |
| Dina baby | 6.91±0.34 | 24.35±0.26* | 8.6±1.27* | 13.01±1.07 | 53.48±0.22* | 3.25±0.21 | 485.11 |
| Tanty | 9.62±0.14* | 19.69±0.0* | 1.0±0.00 | 28.38±0.7 | 36.25±0.64* | 6.75±0.21 | 435.73 |
| Bio nutria | 6.58±0.16 | 11.56±0.29* | 14.2±1.41* | 17.33±1.24 | 58.65±0.63* | 2.25±0.35 | 407.96 |
| Industrial CF | | | | | | | |
| Phosphatine | 4.93±1.73 | 5.0±0.24 | 2.7±0.42* | 24.34±0.01 | 61.80±0.85 | 4.70±0.14 | 389.56 |
| Blesolac | 3.91±0.53 | 12.91±0.26* | 33.1±0.28* | 17.32±1.25 | 64.80±0.85* | 3.85±0.21 | 444.67 |

In the same column, values having the * are significantly different at $p > 0.05$ from that of Phosphatine; CF= Complementary food

Table 2: Vitamin analysis of complementary foods

| Vitamins | Vit C (mg) | β -carotene (μ g) |
|-------------------------------|------------|------------------------------|
| FAO/WHO feeding pattern, 2006 | 15mg | 300UI |
| Semi-scaled CF | | |
| Soyaconia | 0.13±0.40* | 0.00±0.00* |
| Tantyreine | 0.11±0.08* | 17.7±0.63* |
| Banana custard | 0.09±0.01* | 0.00±0.00* |
| Sweet potato custard | 0.10±0.02* | 20.32±0.97* |
| Dina baby | 0.17±0.00* | 13.19±1.12* |
| Tanty | 0.01±0.00* | 14.30±1.31* |
| Bio nutria | 0.21±0.02* | 0.00±0.00* |
| Industrial CF | | |
| Phosphatine | 0.09±0.04* | 0.00±0.00* |
| Blesolac | 0.07±0.00* | 0.00±0.00* |

Values with the * in the column are significantly different from that of Phosphatine; CF= Complementary food

Proximate Analysis

The results (Table 1) showed that the highest moisture content is seen with Tanty (9.62%) while Blesolac recorded the least among (3.91%). The moisture content of Bionutri, Dina Baby, Blesolac, and Soyaconia samples were not significantly ($P > 0.05$) different from that of the standard (Phosphatine); while Banana Custard and Tanty, were significantly ($P < 0.05$) higher from that of the standard. On the same line, Sweet Potato Custard and Tantyreine were also greater ($P < 0.05$) than that of the standard.

The results from the analysis showed that the sample with the highest fat content was Dina baby (24.35%) with the least recorded for Phosphatine (5.0%). When compared with the standard, all the CF samples were significantly different ($p > 0.05$) from that of the standard.

The results for fibre content of the complementary foods indicated that Blesolac had the highest value (33.1%) and Tanty with the least value (1.00%). When compared with the standard (Phosphatine), Tanty was not significantly different ($p>0.05$) but all the other samples were significantly higher than that of the standard (Phosphatine).

Results for protein content indicated that Tanty (28.38%) had the highest with the lowest value recorded for Tantyreine (12.08%). The protein contents for Bionutri, Dina baby, Blesolac, Tanty and Soyaconia were not significant when compared with the standard (phosphatine), meanwhile the remaining seven samples had a significantly lower value when compared with standard.

Carbohydrate content evaluated revealed that Blesolac had the highest value (64.80%) and tanty recorded the least value (36.25%) Banana Custard and Sweet Potato Custard showed no significant difference ($p>0.05$) when compared with standard (Phosphatine). Bionutri, and Soyaconia were significantly lower with the standard with Blesolac being significantly higher ($p<0.05$), whereas the carbohydrate content of all remaining samples was significantly lower ($p<0.05$) from the standard. Tantyreine, Dina baby, Bionutri, Blesolac, and Tanty had lower ($p<0.05$) carbohydrates content which were significantly lower in all samples except sweet potato custard and Blesolac when compared to the carbohydrates content recommended by FAO/OMS (2006) [3].

Ash content for the different complementary foods were recorded in which Tanty had the highest value (6.75%) with Bionutri (2.2%) having the least value. The ash content of the remaining samples was not significantly different at $p>0.05$.

Energy content of the different complementary food samples revealed that all the semi-scaled complementary foods except for sweet potato custard were higher in energy than the standard. The caloric contents of local weaning energy content of the samples which is higher than the recommended value (400 Kcal100g⁻¹) by FOA/OMS Dina Baby (485.11 Kcal100g⁻¹), Soyaconia (420.55Kcal100g⁻¹), Tanty (435.73 Kcal100g⁻¹), Tantyreine (461.22 Kcal/100g), Bio Nutri (407.96 Kcal100g⁻¹) and Blesolac (444.67 Kcal100g⁻¹).

Vitamin Analysis

The vitamin C content of the complementary food samples revealed that Bionutri (0.21mg) and the least value for Tanty (0.01mg). Vitamin C content the remaining samples all showed non-significant difference from the standard ($p>0.05$) for Vitamin C content.

The results for β carotene content were also recorded in table 2, with the highest value recorded for Sweet Potato Custard (20.32 μ g) with the least value (0.002 μ g) for Soyaconia, Banana Custard, Blesolac, Phosphatine, Bionutri, being non- significantly different from that of the standard (phosphatine) at $p>0.05$ with the other samples being significantly higher than the standard (Phosphatine) at $p<0.01$. The results obtained for β carotene (ranging from 0.00 - 20.32 μ g) and C (ranging from 0.01-0.21mg).

Mineral Composition of the Complementary Foods

Mineral content of the complementary foods is presented in Table 3. Potassium(K) and calcium (Ca) were the highest minerals present and ranged from 113.78mg (Banana Custard) to 674.43mg (Soyaconia) and 160(Banana Custard) to 2720mg (Dina baby) respectively. The other minerals and their contents were: Magnesium 19.44mg (banana custard) to 99.63mg (Tantyreine), Zinc 3.84mg (banana custard) to 13.08mg (Phosphatine), Sodium 38.73mg (Banana Custard) to 117.9mg (Soyaconia), Iron 0.09mg (Banana Custard) to 47.90mg (Tantyreine). Iron and Zinc had the lowest values in these complementary foods The mineral analysis results are presented in (Table 3). The amount of sodium, potassium and zinc of the sample Bionutri (325.13-400.89mg) were in the same range as that of Phosphatine standard but all the complementary food samples were not comparable to the sodium, potassium and zinc levels of the infant foods recommended by FAO/WHO (2006). The samples analyzed did not also fall within the recommendation for calcium, zinc and iron, with only one of them being close (Tanty) to the recommended magnesium level. Potassium levels (113.78- 925.27mg).

Table 3: Mineral analysis of complementary foods

| Minerals | Ca mg/100g | Mg mg/100g | Zn mg/100g | Na mg/100g | K mg/100g | Fe mg/100g |
|-----------------------------------|---------------|---------------|---------------|---------------|--------------|---------------|
| FAO/WHO recommended pattern, 2006 | 341.2 | 48.7 | 3.7 | 60 | 408.7 | 8.5 |
| Semi-scaled CF | | | | | | |
| Soyaconia | 1120 | 41.31 | 5.41 | 117.9 | 674.43 | 6.74 |
| Tantyreine | 160 | 99.63 | 6.54 | 60.73 | 358.13 | 47.9 |
| Banana custard | 160 | 53.46 | 3.84 | 38.73 | 113.78 | 0.3 |
| Sweet potato custard | 800 | 43.74 | 5.72 | 60.73 | 358.13 | 14.37 |
| Dina baby | 2720 | 58.32 | 6.96 | 87.12 | 674.43 | 12.83 |
| Tanty | 480 | 48.6 | 7.4 | 60.73 | 358.13 | 0.72 |
| Bionutri | 3360 | 19.44 | 5.51 | 87.12 | 455.57 | 0.09 |
| Industrial CF | | | | | | |
| Phosphatine | 960 | 36.45 | 13.08 | 87.12 | 455.57 | 3.8 |
| Blesolac | 3040 | 43.74 | 5.99 | 60.73 | 268.69 | 0.44 |

CF= Complementary food

The magnesium contents of the flours analyzed (36.45-99.63 mg). The Mg content is high compared to the standard Phosphatine with a value of 36.45mg.

The sodium contents of the flours analyzed (38.73-117.9mg) are higher than that recommended for children by FAO/WHO (2006).

The Fe contents of the flours tested (0.09-14.47mg) are higher as compared to the recommendation of 8.50mg/100g by FAO/WHO (2006).

Microbial Evaluation of the Complementary Foods

The results of microbial analysis were recorded in table 4, with Dina baby recording the highest (2) coliform count at dilution factor 10^{-1} . Banana Custard and Sweet Potato Custard recorded the highest (108 and 40) bacterial count at a dilution of 10^{-2} . Phosphatine showed the highest (17) yeast count at a dilution of 10^{-3} . The results of microbial evaluation (Table 4).

Table 4: Mineral analysis of complementary foods

| Sample | Coliform count | | Total bacterial count | | Yeast count | |
|----------------------|----------------|-----------|-----------------------|-----------|-------------|-----------|
| | 10^{-1} | 10^{-2} | 10^{-2} | 10^{-3} | 10^{-1} | 10^{-2} |
| Dilutions | | | | | | |
| Semi-scaled CF | | | | | | |
| Soyaconia | 0 | 4 | | | | |
| Tanty reine | 0 | 2 | | | | |
| Banana custard | 0 | 108 | | | | |
| Sweet potato custard | 0 | 40 | | | | |
| Dina baby | 2 | | | 15 | | |
| Tanty | 1 | 1 | | | 26 | |
| Bionutri | 0 | 6 | | | 27 | |
| Industrial CF | | | | | | |
| Phosphatine | 0 | 72 | | | | 17 |
| Blesolac | 0 | 1 | | | | |

CF= Complementary food

Functional Properties of the Complementary Foods

The results of functional properties were presented in Table 5. Foam capacity revealed highest value for Phosphatine (3.3%) with lowest value for soyaconia and banana custard (2.05%). The foam capacity of Soyaconia, Tantyreine, Sweet Potatao Custard, Dina baby, Tanty and Bionutri were significantly lower than that of phosphatine.

Table 5: Functional properties of complementary foods

| Samples | Foam capacity (%) | Dispersibility (%) | Loose bulk density (g/ml) | Package density (g/ml) | Water absorption capacity (ml/g) | Swelling capacity (ml/g) | OAC (ml/g) | Foam stability (%) |
|-----------------------|-------------------|--------------------|---------------------------|------------------------|----------------------------------|--------------------------|------------|--------------------|
| Semi-scaled CF | | | | | | | | |
| Soyaconia | 2.05±0.07* | 0.07±0.001* | 0.88±0.00* | 0.64±0.01* | 1.45±0.07* | 1.07±0.00* | 2.05±0.07* | 0.02±0.00* |
| Tantyreine | 2.1±0.14* | 0.07±0.001* | 0.87±0.00* | 0.67±0.00* | 1.05±0.07* | 1.04±0.00* | 2.1±0.14* | 0.02±0.00* |
| Banana custard | 2.05±0.07 | 0.08±0.001* | 1.01±0.01* | 0.69±0.06* | 0.6±0.14* | 1.02±0.03* | 2.05±0.07* | 0.00±0.00 |
| Sweet potato custard | 2.45±0.07* | 0.06±0.001* | 1.07±0.00* | 0.63±0.00* | 2.05±0.07* | 1.20±0.00 | 2.45±0.07* | 0.01±0.00* |
| Dina baby | 3.05±0.07* | 0.07±0.001* | 0.94±0.02* | 0.63±0.02* | 0.55±0.07* | 1.13±0.05 | 3.05±0.07 | 0.02±0.00* |
| Tanty | 3.05±0.07* | 0.07±0.001* | 0.78±0.00* | 0.46±0.01 | 1.2±0.14* | 1.02±0.02* | 3.05±0.07 | 0.04±0.00* |
| Bionutri | 2.1±0.14* | 0.07±0.001* | 1.01±0.00* | 0.73±0.00* | 1.05±0.07* | 1.59±0.01* | 2.1±0.14* | 0.01±0.00* |
| Industrial CF | | | | | | | | |
| Phosphatine | 3.3±0.42* | 0.01±0.004 | 0.72±0.00 | 0.42±0.01 | 3.55±0.07 | 1.15±0.00 | 3.3±0.42 | 0.00±0.00 |
| Blesolac | 2.45±0.07* | 0.07±0.001* | 1.12±0.01* | 0.66±0.06* | 1.05±0.07* | 1.38±0.02* | 2.45±0.07* | 0.01±0.00* |

In the same column, values having the * are significantly different at $p > 0.05$ from that of Phosphatine; CF= Complementary food

Dispersibility was highest in Banana Custard (0.08%) and lowest in Phosphatine (0.01%). The difference between the dispersibility results of the samples compared to that of the standard (Phosphatine) was significantly high at $p < 0.05$. The results of foam capacity recorded that the highest value for Phosphatine (3.32%) with the lowest value for Soyaconia and Banana custard (2.05%). Apart from Dina Baby and Tanty, the others had significantly lower foam capacity as compared to the standard (Phosphatine) at $P > 0.05$.

Loose bulk density results were highest for Soyabeans Pap (1.32g/ml) meanwhile the lowest value was recorded for Bionutri and Banana Custard (0.01g/ml). The results showed that loose bulk density of all the samples were significantly higher as compared to the standard (Phosphatine) at $p < 0.05$. Packaged bulk density results revealed that Bio Nutri had the highest value (0.73g/ml) with Phosphatine (0.42g/ml) recording the lowest score. From the results it is seen that only Tanty is not significantly different from the standard (Phosphatine) at $p > 0.05$ meanwhile packaged bulk density of all the other samples had a high significant difference as compared to the standard.

The result of foam stability indicated that Tanty (0.04%) had the highest score whereas Banana custard and Phosphatine recorded none. The results indicated that the foam stability of Blesolac had a significantly higher value than that of the standard (Phosphatine) at $p < 0.05$, whereas that of Bionutri, Dina Baby, Sweet Potato Custard, Tanty, Tantyreine and Soyaconia had a high significantly higher value when compared with

the standard at $p < 0.05$. The foam stability of the remaining samples showed no significant difference with standard.

The ability of the flours to absorb water (WAC) (Table 5) revealed the highest score recorded by Phosphatine (3.55ml/g) with the lowest scores revealed by Dina Baby (0.55ml/g). From the results, WAC for all the samples showed a significantly ($p < 0.05$) lower value when compared with standard.

Oil absorption capacity (OAC) recorded the highest score (3.3ml/g) for Phosphatine with the lowest (2.05ml/g) for Soyaconia and banana custard, it is revealed that the OAC of Dina Baby and Tanty had no significant difference when compared with the standard (Phosphatine) at $p > 0.05$ meanwhile the remaining complementary foods showed high significant difference with standard at $p < 0.05$.

Protein Solubility of the Complementary Foods

Protein solubilities of the complementary foods are summarized in figure 1 and 2. The commercially sold complementary foods figure 1 shown that Blesolac having two peaks has two proteins with isoelectric PH of 5 and 8 respectively with Phoshatine which showed 3 bands standing for 3 proteins with isoelectric pH of 4, 7 and 9.




Figure 1: Protein solubility result for commercial complementary foods




Figure 2: Protein solubility result for semi scaled complementary foods

The locally produced complementary foods also revealed different protein peaks as seen in figure 2 revealed that Tanytreine had three peaks (three proteins) with isoelectric pH at 2, 4 and 7. Bionutri had two peaks (two proteins) with isoelectric PH of 2 and 5. Soyaconia showed three peaks (three proteins) with isoelectric pH of 4, 7 and 9. Banana custard had just one peak (one protein) with isoelectric pH of 8. Sweet potato custard had three peaks (three proteins) with isoelectric pH of 4 and 8. Tanty revealed two peaks with isoelectric pH of 3 and 8. Lastly Dina baby presented two peaks with isoelectric pH of 2, 7 and 9. Those with three peaks (Phosphatine, Soyaconia, Tanytreine Sweet Potato Custard) are the second best and lastly, those with two peaks (Blesolac, Tanty and Dina Baby) are the least recommended.

Discussions

Low amount of moisture in food samples is best for prolonging the longitivity of food products while high amount of moisture in food samples promotes the growth of microorganisms; thus, it can cause the food to get bad. The amount of moisture in foods is an important factor in determining how prone it is to invasion by microorganisms. The higher the amount of moisture, the higher the risk of spoilage by microorganisms. Low moisture content thus denotes low proliferation of microorganisms [10]. The amount of moisture is used as a quality indicator for complementary foods which is $3-8g \cdot 100g^{-1}$ [11], thus the highest amount of moisture observed in sample Tanty is 9.62%. The fact that the was higher value for moisture in local complementary foods could be due to differences in processing and the preservation.

Fat content of Banana Custard, Sweet Potato Custard, (10.73 and 410.5% respectively) follows what was recommended by Protein Advisory group and the fat content of complementary foods should not exceed 10% (Munasinghe *et al.*, 2013). The amount of fat present in a complementary food has an effect on how long it can last on the shelf. It is due to the fact fat can go through oxidative deterioration, that results in food deterioration. However, increase level of fat mainly saturated fatty acids have been proven to elevate the amount of cholesterol in blood; though, this is not what is observed with unsaturated fats like those found in soy bean and cereals [12].

Locally processed complementary diets with little amount of fiber are so necessary because it aids in the wellbeing of infants, taking note of the urge they have to take in more in order to achieve the energy required daily. The amount of crude fiber for children's diet has to be low (Munasinghe *et al.*, 2013), as diets with high amount of fiber causes poor digestion in children. Thus, complementary foods with low amount of fiber were denoted right as good complementary foods. Dietary fiber content was good when compared to that recommended of FAO/OMS (2006) [3].

The amount of protein was lower in the local formulae apart from Blesolac, Bionutri, Soyaconia and Tanyreine formulae compared to the recommendation of FAO/WHO (2006). The values recorded for protein content were similar to value obtained by Tiencheu *et al.*, (2016) [13]. Proteins are necessary, for quick growth and development of children. The low number of proteins recorded for traditional weaning diets have been a main concern for children's feeding habits. There is a need for formulation as it could act as a feasible way of adding the amount of proteins in local sorghum and maize based complementary diets.

The fact that the carbohydrate content of the complementary foods was in close range to that of the carbohydrates content recommended by FAO/WHO (2006) could be assigned to the fact that most of them are made from carbohydrate sources. According to Munasinghe *et al.*, (2013) the ash content of weaning food should not exceed 5%. Although, in this experimental study, all values are lower but were in acceptable limits as those required by FAO/WHO (2006). This could be a result of the fact that most of these complementary foods are not enhanced with many of mineral ingredients. The caloric contents of most of the complementary diet samples were higher than the required value (400 Kcal100g⁻¹) by FOA/WHO. This high caloric value could be due to their high amount of carbohydrate and also their high fat content as compared to Sweet Potatao Custard and Banana Custard, which had lower caloric values. These results were not in line with that of Asoba *et al.*, (2018) [14]. These results were lower than the results reported by Kanu *et al.*, (2009) [15] who also studied production and evaluation of breakfast cereal-based porridge mixed with sesame and pigeon peas for adults.

Microbiological evaluation was carried out on complementary foods to ascertain their wholesomeness for consumption. In this evaluation, the amount of Coliform was in recommended limits (Table 4) but yeast count and total bacterial load were not. Following the accepted by the UK Food Protection Agency and Food Standard Australia New Zealand (FSANZ), desired Total Plate Count (TPC) for cereal flour mixtures is < 10⁷ CFU/g and coliform count is < 3 CFU/g. The analyzed semi-scaled and commercial complementary diets were in the desired range. The spoilage of many foods will actually take place when the bacteria count reaches 10-100 m/g of product. The international microbiological standard desires a bacteria infestation limit of < 10⁶ CFU.mL⁻¹ for food. The high bacteria load noted in locally made complementary foods like

Banana Custard may be as a result of poor sanitary conditions of vendor and a lack of good production practices during the food processing. These results are different from that of Asoba *et al.*, (2018) [14] who presented lower values for total bacterial count and this could be explained by the fact that our homemade complementary foods could have been self-contaminated by the mothers during preparing either by using dirty hands or dirty kitchen equipment.

The results obtained for β carotene (ranging from 0.00 - 20.32 μ g) and C (ranging from 0.001-0.21mg) are not in line with the higher values observed for complementary foods processed from cooked banana, fortified with cowpea and peanut as observed by Bassey *et al.*, (2013) [16]. These results prove that these complementary diets might not be used as an appropriate source of vitamins especially vitamins A and C which is essential for visual and immune system development in infant and children. These results are different from that of Asoba *et al.*, (2018) [14] who presented higher values and this could be due to the fact that these vitamins are affected by production process as they are not thermo-stable.

Minerals are vital for health. In this study the mineral content in most of the complementary foods including the control phosphatine did not meet the recommendation of FAO/OMS (2006), except for iron, thus they can protect against iron deficiency anemia. Also, the semi-scaled complementary food, Tanty showed good amounts of all the five minerals analyzed except for Fe. Very low amount of iron was observed in all the other complementary food samples and iron deficiency can result in anemia.

For the functional properties, the differences showed in water absorption among the flours may be due to different protein concentration, their degree of interaction with water and their conformational characteristics (McWatters *et al.*, 2003). On the other hand, McWatters *et al.*, (2003) stated that lower water absorption capacity is caused by less availability of polar amino acids in flours. This effect is most likely due to lose interaction of amylose and amylopectin in the native granules of starch and softer interaction enhances maintenance of the granules structure [17]. This result revealed that the composite flours had better gelling property. The ability of a flour to absorb water is necessary in bulking and consistency of product and also in baking applications [18].

Bulk density (loose bulk density and packaged density) presents an indication of the relative volume of packaging material necessary and high bulk density is a better physical factor when evaluating the mixing quality of a particulate matter Achinewhu *et al.*, (1998) [19]. The bulk density is a view of the amount the flour samples can take, if left to rest directly on one another. The density of processed products indicates the characteristics of its container or package product density indicates the number and strength of packaging material, texture or mouth feel (Lewis,1992). According to Basman *et al.*, (2003) [20], higher bulk density is needed for greater ease of dispersibility of flours. Also, high bulk density reduces the caloric and amount of nutrients per feed of children which can cause poor growth. However, low bulk density would be a good factor in the formulation of complementary foods (Ugwu and Ukpabi, 2002). The low bulk density values observed could thus be good in the preparation of complementary food formulas.

Oil absorption capacity (OAC) has been denoted to be due to physical entrapment of oil and the binding of fat to the polar chains of proteins. Kinsella (1976) [7] observed that more water hating proteins show greater binding of lipids, this indicates that non-polar amino acids side chains bind the paraffin chains of fats. The

lower OAC of the formulated blends (Soyaconia and Banana custard) could be as a result of the fact that Phosphatine (with a greater OAC) had more available non-polar side chains in its protein molecules than Soyaconia, Banana custard and Bionutri. This implies that Phosphatine will be able to hold more flavour and most likely have a nicer mouth feel as oppose to the other complementary diets.

Swelling power refers to the expansion accompanying spontaneous uptake of solvent while solubility index (SI) is the number of water-soluble solids per unit weight of the sample. According to Kinsella (1976) [7], Swelling causes changes in hydrodynamic properties of the food thus impacting characteristics such as body, thickening and increase in viscosity to foods.

Protein solubility denotes the number of soluble proteins present in a sample and also their associated isoelectric points. Therefore, a sample with the highest number of peaks denotes more soluble proteins with the corresponding troughs revealing the associated isoelectric points of the proteins [21-29].

Conclusion

When nutritional evaluation of these local weaning foods was carried out, it was observed that amount of moisture in Phosphatine was within the required limits. The amount of carbohydrate in the complementary food Sweet Potato Custard was in the acceptable limit. Potassium and calcium were the highest minerals present in the local complementary foods with one of the samples Tanty having the acceptable value for Magnesium mineral. The amount of both vitamins which were analyzed were very low. The functional properties of the complementary diets showed a significant difference when compared to that of Phosphatine. It was also observed during the microbiological evaluation that, the complementary foods analyzed fell in the accepted limits of food protection free from bacteria and fungi. However, the bacteria load was a little high which could have resulted from sanitary conditions. Thus, it could be seen that most of the complementary diets are inadequate in or more nutrients and need fortification in order to serve as a good complementary food for children.

Acknowledgements

We are grateful to the laboratory of Nutrition and Nutritional Biochemistry, Department of Biochemistry, University of Yaounde 1 for supporting this study through our supervisor/ Mentors; Prof Julius Oben and Prof. Azantsa Kingue. Authors are also thankful to Islamic Development Bank Postdoctoral Programme (IsDB) for their financial life stipend in support of this research work granted to Dr. Tiencheu Bernard. We are also very grateful to mothers that give their consent for data collection to undertake this study.

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Bibliography

1. AOAC (2005). Official methods of analysis of the association of official analytical chemists international.
2. Adesetan, T. O., Egberongbe, H. O., Illusanya, O, A. & Olurunjuwon, O. (2014). Antimicrobial Sensitivity of bacterial isolates from street vended fruits in Ijebu area of Ogun state, Nigeria. *International Research Journal of microbiology*, 4(9), 220-225.
3. FAO/OMS. (2006). Programme mixte FAO/OMS sur les normes alimentaires. Rapport des vingt-septième sessions du comité du codex sur la nutrition et les aliments diététiques ou de régime. ALINOM 06/29/26,105p. Randomized trials in Honduras. *J Nutr.*, 131, 262-267.
4. Pierro, J., Abulaimoun, B., Roth, O. & Blau, J. (2016). Factors associated with supplemental formula feeding of breastfeeding infants during post-partum hospital stay. *Breastfeed Med.*, 11, 196-202.
5. Agunod, M., Yamaguchi, N., Lopez, R., Luhby, A. & Glass, G. (1969). Correlation study of hydrochloric acid, pepsin and intrinsic factor secretion in newborns and infants. *Dig Dis Sci.*, 14(6), 400-414.
6. Pullen, D., Kennedy, G. E. (2005). From the ape's dilemma to the weanling's dilemma: early weaning and its evolutionary context. *J Hum Evol.*, 48(2), 123-145.
7. Kinsella, J. E. (1976). Functional properties of proteins in foods: A survey. *Critical Reviews in Food Science & Nutrition*, 7(3), 219-232.
8. Kulkarni, K., Kulkarni, D. & Inglee, U. (1991). Sorghum malt-based weaning formulations: Preparation, functional properties and nutritive value. *Food and Nutrition Bulletin*, 13(4), 1-7.
9. Naranyanga, k. & Narasinga, M. (1982). Functional properties of raw and heat processed winged bean (*Psophocarpus tetragonolobus*) flour. *J. Sci.*, 47(5), 1534-1538.
10. Achidi, A. U., Moyeh, M. N., Tiencheu, B., Tenyang, N., Womeni, H. M., Ebini, L. T. & Tatsinkou, F. (2016). Quality evaluation of nine instant weaning foods formulated from cereal, legume, tuber, vegetable and crayfish. *International Journal of Food Science and Nutrition Engineering*, 6(2), 21-31.
11. Amankwah, E. A., Barimah J., Nuamah, A. K. M., Oldham, J. H. & Nnaji, C. O. (2009). Formulation of weaning food from fermented maize, rice, soybean & fish meal. *Pakistan Journal of Nutrition*, 8, 1747-1752.
12. Szalai, G., Janda, T., Páldi E. & Dubacq, J. (2001). Changes in the fatty acid unsaturation after hardening in wheat chromosome substitution lines with different cold tolerance. *J. Plant Physiol.*, 158(5), 663-666.
13. Tiencheu, B., Tjepma, E. F., Achidi, U., Tatsinkou, F. B. & Tenyang, W. (2016). Formulation and nutritional evaluation of instant weaning foods processed from maize (*Zea mays*), pawpaw (*Carica papaya*), red beans (*Phaseolus vulgaris*) and mackerel fish meal (*Scomber scombrus*). *American Journal of Food Science and Technology*, 4(5), 149-159.

14. Asoba, G. N., Sumbele, I. N., Anchang, K. J., Samuel, M. & Kaptso, K. G. (2018) Nutritional evaluation of commonly used local weaning food processed and sold in the Mount Cameroon Region. *International Journal of Food Science and Nutrition Engineering*, 8(6), 131-141.
15. Kanu, P., Sandy, E., Kandeh, J., *et al.* (2009). Production and evaluation of breakfast cereal-based porridge mixed with sesame and pigeon peas for adults. *Pakistan Journal of Nutrition*, 8(9), 1335-1343.
16. Bassey, F. I., Mcwatters, K. H., Edem, C. A. & Iwegbue, C. M. A. (2013). Formulation and nutritional evaluation of weaning food processed from cooking banana, supplemented with cowpea and peanut. *Food Science & Nutrition*, 1(5), 384-391.
17. Lorenz, K. & Collins, F. (1990). Quinoa (*Chenopodium quinoa*), Starch - physicochemical properties and functional characteristics. *Starch/starke*, 42(3), 81-86.
18. Niba, L. L., Bokanga, M., Jackson, F. I., Schlimme, D. S. & Li, B. W. (2001). Physicochemical properties and starch granular characteristics of flour from various *Manihot esculenta* (cassava), genotype. *J. Ed. Sci.*, 67(5), 1701-1705.
19. Achinewhu, S. C., Barber, L. I. & Ijeoma, I. O. (1998). Physicochemical properties and garrification of selected cassava cultivars in River State. *Plant Food Hum. Nutr.*, 52(2), 133-140.
20. Basman, A., Koksel, H. & Ng, P. K. (2003). Utilization of Transglucanase use to increase the level of barley and soy flour incorporation in wheat flour breads. *J. Food Sci.*, 68(8), 2453-2460.
21. Anigo, K. M., Ameh D. A., Ibrahim S. & Danbauchi S. S. (2010). Nutrient composition of complementary food gruels formulated from malted cereals, soybeans and groundnut for use in North-western Nigeria”, *African Journal of Food Science*, 4(3), 65-72.
22. FAO (2001). *Improving nutrition through home gardening*. A training package for preparing field workers in Africa. FAO Rome.
23. Egonlety, M. (2002). Production of legume-fortified weaning food. *Food Res. Inter.*, 35(2-3), 233-237.
24. Kanu, P. J., Kerui, Z., Huiming, Z., Haifeng, Q., Kanu, J. B. & Kexue, Z. (2007a). Sesame protein 11: Functional properties of sesame (*Sesamum indicum* L.) protein isolate as influenced by pH, temperature, time and ratio of flour to water during its production. *Asian J. Biochem.*, 2, 289-301.
25. Lewis, M. J. (1990). *Physical properties of Food and Food Processing Systems*. Hartnolls Limited, Bodman Cornwall, Great Britain.
26. Ogonnaya, J. A., Ketiku, A. O., Mojekwu, C. N., Mojekwu, J. N. & Ogonnaya J. A. (2012). Energy, iron and zinc densities of commonly consumed traditional complementary foods in Nigeria. *British Journal of Applied Science and Technology*, 2(1), 48-57.

-
27. Ramakrishnan, U., Martorell, R., Schroeder, D. & Flores R. (1999). Role of intergenerational effects on linear growth. *Journal of Nutrition*, 129(2S Suppl), 544S-549.
28. Reyes, H., Perez-Cuevas, R., Sandoval, A., Castillo, R., Santos, J. I., *et al.* (2004). The family as a determinant of stunting in children living in extreme poverty: a case Control study. *BMC Public Health.*, 4, 57.
29. Tenagashaw, M. W., Kenji, G. M., Melaku, E. T., *et al.* (2016). Proximate composition and selected functional properties of complementary foods from teff fortified with soybean and orange-fleshed sweet potato. *RUFORUM Working Document Series No. 14*(1), 953-965.